

3 in

**Drug Facts**


**Active ingredients Purpose**  
Benzocaine 5%.....Oral Anesthetic  
Glycerin 33%.....Oral Demulcent

**Uses** for temporary relief of pain associated with canker sores.

**Warnings**  
Do not exceed recommended dosage.  
**Methemoglobinemia warnings:**  
use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood.

**Allergy alert** • do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

**Do not use** this product for more than 7 days unless directed by a doctor. If irritation, pain or redness persists or



**EFFICIENT®**

**FINAFTA®**

Benzocaine 5%  
Glycerin 33%

**Anesthetic/Analgesic Liquid**

**Instant Relief for the Mouth and Gums**

**0.5 FL. Oz. (15 mL)**

**Drug Facts (continued)**

worsens; or if swelling, rash or fever develops, see your doctor promptly.

**Keep out of reach of children.**

**Directions**  
•adults and children over 12 years of age apply to the affected area, 4 times daily. Do not rinse mouth after each use •children between 2 and 12 years of age: consult a doctor and supervise in the use of the product  
•do not use in children under 2 years of age.


**Inactive ingredients**  
menthol, methylparaben, polyethylene glycol, propylparaben, propylene glycol, purified water, phosphoric acid, sucralose.

**Keep the box for complete Drug Facts information in English and Spanish.**

Distributed by: Efficient Laboratories, Inc.®  
Miami, FL 33166,  
Made in U.S.A.

1 1/2 in

3 in



1 1/2 in