

5.5 Inch

Drug Facts

Active ingredients (in each 30 mL dose cup)

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

Purposes

Uses

temporarily relieves these common cold/flu symptoms:

- minor aches and pains • headache • sore throat • runny nose and sneezing • cough • relieves cough to help you get to sleep • temporarily reduces fever.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 6 doses in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks daily while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash.

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.


• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

• to make a child sleep.

Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • glaucoma • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema • a sodium-restricted diet.

Ask a doctor or pharmacist before use if you are • taking the blood thinning drug warfarin • taking sedatives or tranquilizers.

When using this product • do not use more than directed • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • alcohol, sedatives, and tranquilizers may increase drowsiness • be careful when driving a motor vehicle or operating machinery •

EFFICIENT®

Rompe Pecho®
NT

Nighttime Severe
Cold & Flu Relief

Pain Reliever/Fever Reducer (Acetaminophen)
Cough Suppressant (Dextromethorphan HBr)
Antihistamine (Doxylamine Succinate)

ADVANCED FORMULA
ALCOHOL FREE

Adults & Children 12 years & over

Contents: 6 FL Oz (180 mL)

Drug Facts (continued)

Stop use and ask a doctor if • nervousness, dizziness, or sleeplessness occur • pain or cough gets worse or lasts more than 7 days • new symptoms occur • symptoms occur • fever gets worse or lasts more than 3 days • redness or swelling is present • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

If pregnant or breast-feeding, ask a health professional before use.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not take more than directed (see overdose warning) • do not exceed more than 6 doses in any 24 hours period • shake well before use • measure only with dosing cup provided • keep dosing cup with product • mL = milliliter • TBSP = tablespoonsful • this adult product is not intended for use in children under 12 years of age • adults and children 12 years of age and over: 30 mL every 4 hours • children under 12 years of age: do not use.

Other information • each 30 mL contains: potassium 20 mg; sodium 12 mg • store between 15-30°C (59-86°F). Do not refrigerate • product color & flavor may vary due to natural honey.

TAMPER-EVIDENT FEATURE: Do not use this product if printed safety seal is torn, broken, or missing.

Inactive ingredients blue cohosh root extract, citric acid, echinacea root extract, eucalyptus oil, flavor, ginkgo biloba leaf extract, glycerin, goldenseal root extract, honey, horsehound (flower, leaf, stem) extract, licorice root extract, menthol, methylparaben, mullein leaf extract, myrrh gum, potassium citrate, potassium sorbate, propylene glycol, propylparaben, slippery elm bark extract, sodium chloride, sucralose, water, wild cherry bark extract and zinc sulfate.

Questions or Comments? 305-805-3456
Monday - Friday (9 AM - 5 PM EST) or www.efficientlabs.com

Other cold formulas: Rompe Pecho® DM, Rompe Pecho® SF, Rompe Pecho® EX, Rompe Pecho® CF, Rompe Pecho® FLU, Rompe Pecho® MAX, Rompe Pecho® Cough Quilt®, Melts y Miel®.

Find the product brochure in the box to obtain information in Spanish.
Keep the box for complete instructions.

Rev. 1: 10/2023

Distributed by:
Efficient Laboratories, Inc.®
Miami, FL 33166, Made in U.S.A.

3 Inch