

6.7 Inch.

Drug Facts

Active ingredients (in each 20 mL) Purposes

Acetaminophen, 650 mg.....	Pain reliever/fever reducer
Dextromethorphan, HBr 20 mg.....	Cough suppressant
Guaifenesin, 400 mg.....	Expectorant
Phenylephrine HCl, 10 mg.....	Nasal decongestant

Uses - temporarily relieves these common cold and flu symptoms:
 - nasal congestion - sinus congestion and pressure - cough due to minor throat and bronchial irritation - minor aches and pains - sore throat - headache
 • temporarily reduces fever - promotes nasal and/or sinus drainage
 • temporarily relieves sinus congestion and pressure - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings
Liver warning: this product contains acetaminophen.
Severe liver damage may occur if: - you take more than 6 doses in 24 hours, which is the maximum daily amount for this product - taken with other drugs containing acetaminophen - 3 or more alcoholic drinks every day while using this product.
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: - skin reddening - blisters - rash. If skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use - with any other drug containing acetaminophen (prescription or nonprescription) - if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist - if you have persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema, or where cough is accompanied by excessive phlegm (mucus) unless directed by a doctor - if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
 • heart disease • liver disease • high blood pressure • thyroid disease • diabetes • difficulty in urination due to an enlarged prostate gland • a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema • a cough that is followed by excessive phlegm (mucus).

11/2022 REV.1 **Keep box for complete instructions**

EFFICIENT

Rompe Pecho®

MAX Multi Symptoms

**PAIN RELIEVER
 FEVER REDUCER (Acetaminophen)
 COUGH SUPPRESSANT
 (Dextromethorphan HBr)
 EXPECTORANT (Guaifenesin)
 NASAL DECONGESTANT
 (Phenylephrine HCl)**

Maximum Strength

Adults & Children
 12 years & over

NET WEIGHT: 8 FL OZ (237 mL)

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are
 - taking the blood thinning drug warfarin - taking tranquilizers or sedatives.

When using this product do not use more than directed

Stop use and ask a doctor if
 • symptoms do not improve within 7 days, tend to recur, or are accompanied by a fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition - nervousness, dizziness or sleeplessness occurs - pain and nasal congestion gets worse or last more than 5 days (children) or 7 days (adults) - fever gets worse or last more than 3 days - redness or swelling is present or new symptoms occur.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions
 • do not exceed more than 6 doses in any 24 hours period - shake well before use - measure only with dosing cup provided - keep dosing cup with product - mL = milliliter - this adult product is not intended for use in children under 12 years of age - adults and children 12 years of age and over: 20 mL every 4 hours - children under 12 years of age: do not use.

Other information - each (20 mL) contains: potassium 15 mg, sodium 8 mg - product color & flavor may vary due to natural honey - store between 15-30°C (59-86°F) - do not refrigerate.

Tamper-Evident Feature: Do not use this product if printed safety seal is torn, broken, or missing.

Inactive ingredients blue cohosh root extract, citric acid, echinacea root extract, eucalyptus oil, ginkgo biloba leaf extract, glycerin, goldenseal root extract, honey, licorice root extract, licorice root extract, menthol, methylparaben, mullein leaf extract, myrrh gum, potassium citrate, potassium sorbate, propylene glycol, propylparaben, slippery elm bark extract, sodium chloride, sucralose, water, wild cherry bark extract and zinc sulfate.

Questions or comments? 305-805-3456 Monday to Friday from 9 am to 5 pm, EST or www.efficientlabs.com

Other cold formulas: Rompe Pecho® EX, Rompe Pecho® SF, Rompe Pecho® CF, Rompe Pecho® DM, Cough Out® and Mefim y Mas®, Rompe Pecho® FLU and Rompe Pecho® NT.

Distributed by:
Efficient Laboratories, Inc.
 Miami, FL 33166, Made in U.S.A.

3.07 Inch.